



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

VACCINE-INDUCED MASSIVE PULMONARY EMBOLISM AND
THROMBOCYTOPENIA FOLLOWING A SINGLE DOSE OF
JANSSEN AD26.COVID.S VACCINATION

Rosa Curcio , Vito Gandolfo , Riccardo Alcidì ,
Luciano Giacomino , Tommaso Campanella , Genni Casarola ,
Rachele Rossi , Lorenzo Chiatti , Marco D'abbondanza ,
Rita Commissari , Paolo Gresele , Giacomo Pucci ,
Gaetano Vaudo

PII: S1201-9712(21)01244-3
DOI: <https://doi.org/10.1016/j.ijid.2021.12.345>
Reference: IJID 5902

To appear in: *International Journal of Infectious Diseases*

Received date: 16 August 2021
Revised date: 1 December 2021
Accepted date: 18 December 2021

Please cite this article as: Rosa Curcio , Vito Gandolfo , Riccardo Alcidì , Luciano Giacomino , Tommaso Campanella , Genni Casarola , Rachele Rossi , Lorenzo Chiatti , Marco D'abbondanza , Rita Commissari , Paolo Gresele , Giacomo Pucci , Gaetano Vaudo , VACCINE-INDUCED MASSIVE PULMONARY EMBOLISM AND THROMBOCYTOPENIA FOLLOWING A SINGLE DOSE OF JANSSEN AD26.COVID.S VACCINATION, *International Journal of Infectious Diseases* (2022), doi: <https://doi.org/10.1016/j.ijid.2021.12.345>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 The Author(s). Published by Elsevier Ltd on behalf of International Society for Infectious Diseases.

This is an open access article under the CC BY-NC-ND license
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

HIGHLIGHTS

- Thrombosis and thrombocytopenia after Sars-CoV-2 viral vector vaccine suggests VITT.
- Thrombosis in unusual sites (splenic, portal, mesenteric, adrenal) is typical of VITT.
- Anti-PF4 antibody tests, if available, should be rapidly determined for diagnosis.
- IVIg could mask functional tests to confirm VITT diagnosis.
- Treatment with IVIg may be associated to false negative functional tests.

**VACCINE-INDUCED MASSIVE PULMONARY EMBOLISM AND THROMBOCYTOPENIA
FOLLOWING A SINGLE DOSE OF JANSSEN AD26.COV2.S VACCINATION**

Rosa CURCIO ^{1,2}
Vito GANDOLFO ^{1,2}
Riccardo ALCIDI ^{1,2}
Luciano GIACOMINO ^{1,3}
Tommaso CAMPANELLA ^{1,2}
Genni CASAROLA ^{1,2}
Rachele ROSSI ^{1,2}
Lorenzo CHIATTI ^{1,2}
Marco D'ABBONDANZA ^{1,2}
Rita COMMISSARI ^{1,3}
Paolo GRESELE ¹
Giacomo PUCCI ^{1,2}
Gaetano VAUDO ^{1,2}

1 Department of Medicine and Surgery, University of Perugia, Perugia, Italy

2 Unit of Internal Medicine, "Santa Maria" Terni University Hospital, Terni, Italy

3 Unit of Emergency, "Santa Maria" Terni University Hospital, Terni, Italy

All authors had access to the data and a role in writing the manuscript.

Potential conflicts of interest: none.

Address for correspondence and request for reprints:

Giacomo Pucci MD, PhD

Department of Medicine and Surgery, University of Perugia,

Unit of Internal Medicine, Terni University Hospital,

Piazzale Tristano di Joannuccio, 1

IT-05100 Terni, Italy

Phone and fax: +39-0744-205201 – E-mail: giacomo.pucci@unipg.it

KEYWORDS

COVID-19, Janssen, AD26.COV2.S, Vaccination, Vaccine-induced thrombosis with thrombocytopenia, pulmonary embolism

ABSTRACT

Vaccine-induced immune thrombotic thrombocytopenia (VITT) has emerged as a rare side effect of adenoviral vector-based vaccines against Coronavirus disease 2019 (COVID-19), and it is most frequently reported after Vaxzevria (AstraZeneca) vaccine. We described a case of severe thrombocytopenia associated with massive pulmonary embolism and portal vein thrombosis, occurring 13 days after the administration of the single dose adeno viral vector-based vaccine Ad26.COV2.S (Janssen vaccines, Leiden, Netherlands). Based on an early clinical suspect, the patient quickly received treatment with corticosteroids and intravenous immunoglobulins, followed by a rapid increase in platelet count that allowed full dose anticoagulation to be timely administered. Treatment with intravenous immunoglobulins, however, could mask the ability of anti-PF4-heparin antibodies to bind and activate platelets in the presence of heparin, leading to false negative results at the immunoassay functional test. Therefore, if VITT is suspected, blood samples for diagnostic confirmation should be collected prior to any treatment to improve diagnostic performance.

CASE REPORT

A 68-years-old man presented to the emergency department for left leg swelling with subacute pain, associated with weakness, dizziness and progressive dyspnoea. His medical history included hypertension and an euthyroid nodular goiter; personal history of thrombosis were negative. Molecular test for SARS-Cov2 nucleic acid (PCR) was negative and anti-spike IgG levels were 316 AU/ml. Vaccination for Coronavirus disease 2019 (COVID-19) with a single dose of Janssen Ad26.COV2.S vaccine (Janssen vaccines,

Leiden, The Netherlands) was reported 13 days before. He denied any previous exposure to anticoagulants including heparins.

The patient was slightly dyspnoic and tachypnoic in room air with peripheral oxygen desaturation, his left leg was swollen and painful. The remaining examination was unremarkable. A point-of-care ultrasound evaluation revealed left femoro-popliteal deep vein thrombosis (DVT) and no signs of right heart overload/dysfunction. A total-body contrast-enhanced computed tomography-scan (CT-scan) showed massive bilateral pulmonary artery embolism and a right intra-hepatic portal thrombosis (Figure 1). Laboratory tests revealed severe thrombocytopenia (platelet count 7000/ μ l), markedly elevated levels of D-dimer (32533 ng/L), increase in serum lactate dehydrogenase (LDH) and C-reactive protein; values of fibrinogen, PT and aPTT were all within the normal range (Table).

Vaccine-induced immune thrombocytopenia (VITT) was suspected, therefore high-dose parenteral corticosteroids (methylprednisolone 80 mg IV daily) and intravenous immunoglobulins (1 g/kg/day for 2 days) were immediately started. An inferior caval vein filter (IVF) was implanted.

High titer antibodies to platelet factor 4 (PF4)–polyanion complexes were identified (optical density (OD) 2.21, negative <0.4) by enzyme-linked immunosorbent assay (ELISA Lifecodes PF4-Immucor-USA), whereas the rapid immunological chemiluminescence assay (HemosIL AcuStar HIT-IgG) was negative, in agreement with the known pattern of VITT-associated antiPF4-antibody positivity. The Heparin-induced platelet aggregation (HIPA) assay was also performed but it resulted negative. Serological tests for human immunodeficiency virus (HIV), hepatitis B and C viruses, TORCH screen, lupus-anticoagulant, anticardiolipin and anti β_2 -glycoprotein antibodies were all negative. Blood levels of C and S proteins, C3 and C4 complement-fractions were within the normal range.

Reduced dose parenteral anticoagulation with fondaparinux was started at day 4, when platelet count reached $20000/\text{mm}^3$ and direct oral anticoagulation was started at day 6 when platelet count exceeded $50000/\text{mm}^3$. At day 16, a Doppler ultrasound showed recanalization of lower limb proximal veins, but IVF retrieval was postponed due to the presence of thrombosis of the caval vein filter. Patient was eventually discharged in good clinical conditions under direct anticoagulants; values of platelets, fibrinogen and LDH had returned in normal ranges.

We report a case of severe thrombocytopenia and venous thromboembolism occurring 13 days after administration of the single dose Janssen COVID-19 vaccine. To the best of our knowledge, this is the first case of VITT associated to the Ad26.COV2.S vaccine described in Italy.

On April 4th, 2021, the European Medicines Agency safety Committee provided a systematic assessment of thromboembolic events associated with thrombocytopenia following the administration of the ChAdOx1 nCoV-19 viral vector-based vaccine (Vaxzevria, AstraZeneca, University of Oxford, and Serum Institute of India). A total of 169 cases of cerebral venous sinus thrombosis (CVST) and 53 cases of splanchnic vein thrombosis were reported to EudraVigilance, leading to the conclusion that a causal relationship between vaccination with Vaxzevria and thromboembolic events with thrombocytopenia was at least a reasonable possibility. Later, given that 6 cases of CVST with thrombocytopenia were also identified in the United States among recipients of approximately 7 million doses of Janssen vaccine, the U.S. Food and Drug Administration and the Centers for Disease Control and Prevention suggested temporary pausing of the administration of Janssen vaccine to allow further investigation [1].

VITT has been mainly reported among females aged <55 years old and between 4 and 16 days after receiving adenoviral vector-based vaccines, although exceptions exist, such as

cases observed in males, in subjects up to 77 years old, and occurring as early as 2 to late 28 days after vaccination [Gresele, 2021a]. Its pathogenesis is still unclear and under investigation, but many reports highlight similarities with autoimmune heparin-induced thrombocytopenia (HIT), tracking a continuum between different platelet-activating anti-PF4/heparin disorders. Autoimmune HIT is a form of HIT without any previous exposure to heparin. A possible role of the adenoviral vector platform and/or of free nucleic acids in the development of the autoimmune response have been suggested [Gresele, 2021a]. *Mc Gonagle* et al. suggested that the local tissue micro-trauma following vaccine inoculation brings adenoviral DNA in contact with platelet-factor 4 (PF4), thus increasing anti-PF4 autoantibody production in susceptible subjects [Mc Gonagle, 2021]. Other hypotheses related the syndrome to the adenoviral vector platform carrying a message from a RNA virus with the consequent random splicing of the DNA message producing unwanted circulating soluble spike protein [Gresele, 2021a].

Venous thrombosis in VITT typically occurs in unusual sites, including cerebral, splanchnic (splenic, portal, mesenteric, adrenal), and ophthalmic veins. Anticoagulation is a cornerstone of HIT-related thrombosis but is prevented by severe thrombocytopenia and DIC. Full-dose anticoagulants should be administered as soon as possible in VITT patients, provided that concurrent thrombocytopenia is corrected [Gresele, 2021b]. To this end IVIg administration, or in severe/refractory cases plasma exchange, are critical to interrupt the immune-mediated mechanisms causing VITT and to obtain a fast increase in platelet count that, in turn, allows full dose anticoagulants to be safely administered in a timely manner [Gresele, 2021b]. In our patient a fast satisfactory response to high-dose corticosteroids and IVIg treatment was obtained in terms of platelet counts allowing full dose anticoagulation to be applied with resolution of the acute clinical picture. Even if there is paucity of data about optimal anticoagulant treatment of VITT, principles of guidance are

based on the use of non-heparin-based anticoagulants (e.g. fondaparinux, argatroban), and to switch to direct oral anticoagulants when the platelet count reaches $50000/\text{mm}^3$ [2]. Two positive assays are conventionally required to confirm a suspected VITT: a quantitative ELISA assay, which detects and quantifies antiPF4 antibodies, and a functional assay, which assesses the ability of anti-PF4-heparin antibodies to bind and activate platelets in the presence of heparin. However, the simultaneous administration of IVIg could mask their functional ability to activate platelets [Bourguignon, 2021]. This might also be of importance because if VITT is suspected, as it was in our case, blood samples for diagnostic confirmation should be collected prior to any treatment to improve diagnostic performance. When treatment has to be started as soon as possible and immunoassay tests are not rapidly available, d-dimer levels may guide the clinical management [Zazzeron, 2021].

Many questions related to VITT remain unsolved: first, what are the viral-vector vaccine components responsible of VITT. Second, what are the molecular mechanisms behind the heterogeneity of results of anti-PF4 functional assays. Third, if it is possible to screen for populations susceptible to VITT and last if vaccinated people with VITT develop more severe symptoms after SARS-CoV-2 re-exposure. Further studies are needed to solve these important and urgent questions.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors

Ethical approval: not required

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

1. <https://www.fda.gov/news-events/press-announcements/fda-and-cdc-lift-recommended-pause-johnson-johnson-janssen-covid-19-vaccine-use-following-thorough>, accessed on 16.08.2021
2. https://cdn.ymaws.com/www.isth.org/resource/resmgr/ISTH_VITT_Guidance_2.pdf, accessed on 16.08.2021
3. Bourguignon A, Arnold DM, Warkentin TE, Smith JW, Pannu T, Shrum JM, et al. Adjunct Immune Globulin for Vaccine-Induced Thrombotic Thrombocytopenia. *N Engl J Med* 2021. doi: 10.1056/NEJMoa2107051. Epub ahead of print.
4. Gresele P, Momi S, Marcucci R, Ramundo F, De Stefano V, Tripodi A. Interactions of adenoviruses with platelets and coagulation and the vaccine-associated autoimmune thrombocytopenia thrombosis syndrome. *Haematologica*. 2021 (2021a). Epub ahead of print. doi: 10.3324/haematol.2021.279289
5. Gresele P, Marietta M, Ageno W, Marcucci R, Contino L, Donadini MP et al. Management of cerebral and splanchnic vein thrombosis associated with thrombocytopenia in subjects previously vaccinated with Vaxzevria (AstraZeneca): a position statement from the Italian Society for the Study of Haemostasis and Thrombosis (SISST). *Blood Transfus* 2021 (2021b); 19(4): 281-3 doi: 10.2450/2021.0117-21. Epub ahead of print.
6. McGonagle D, De Marco G, Bridgwood C. Mechanisms of Immunothrombosis in Vaccine-Induced Thrombotic Thrombocytopenia (VITT) Compared to Natural SARS-CoV-2 Infection. *J Autoimmun* 2021;121:102662. doi: 10.1016/j.jaut.2021.102662.
7. Zazzeron L, Rosovsky RP, Bittner EA, Chang MG. Comparison of Published Guidelines for the Diagnosis and the Management of Vaccine-Induced Immune Thrombotic Thrombocytopenia. *Crit Care Explor*. 2021;3:e0519

Table 1. Laboratory Characteristics of the Index Patient.

Laboratory Analysis	Reference Value	Day 1	Day 4	Discharge
Hemoglobin (g/dl)	13.5 – 18.0	12.2	10.2	11.3
Platelet count (per mm ³)	140000 – 440000	7000	24000	230000
Leukocytes (per mm ³)	4.500 – 10.800	6.350	5.540	8.420
Partial thromboplastin time (sec)	22 - 32	25	28	22
International normalized ratio	0.85 – 1.20	1.08	1.33	1.05
Thrombin time (sec)	70 - 120	86	61	91
Fibrinogen (mg/dl)	160 - 420	217	58	350
D-dimer (mg/liter)	0 – 350	32533	ND	ND
Aspartate aminotransferase (U/liter)	0 - 41	46	33	16
Alanine aminotransferase (U/liter)	3 - 63	60	56	32
γ-Glutamyltransferase (U/liter)	0 - 55	108	112	ND
Lactate dehydrogenase (U/liter)	0 - 248	302	195	ND
C-reactive protein (mg/dl)	0 – 0.8	5.5	ND	ND